

Selvaraj SHALINI^a, Pakshirajan LAKSHMINARASIMHAN^{b,†}, Wilson ARISDASON^c and Debabrata MAITY^{d,*}: **Lectotypification of Two Names in the Genus *Argyreia* (*Convolvulaceae*)**

^aCentral National Herbarium, Botanical Survey of India, Botanic Garden P.O., Howrah, West Bengal, 711103 INDIA;

^bUttam Townscapes, Yerawada, Pune, Maharashtra, 411006 INDIA;

^cBotanical Survey of India, Southern Regional Centre, Lawley Road, Coimbatore, Tamil Nadu, 641 003 INDIA;

^dTaxonomy and Biosystematics Laboratory, Department of Botany, University of Calcutta, 35, Ballygunge Circular Road, Kolkata, West Bengal, 700019 INDIA

*Corresponding author: debmaity@yahoo.com

(Accepted on October 27, 2020)

Summary: Lectotypes of two names in the genus *Argyreia* (*Convolvulaceae*), viz. *A. fulgens* and *A. lanceolata* have been designated in accordance with the International Code of Nomenclature.

Argyreia Lour. is one of the most species-rich genera in the family *Convolvulaceae*. In India, the genus is represented by 44 species and three varieties (Shalini et al. 2017b). Of which, six names in Indian *Argyreia* have been lectotypified in three earlier publications (Shalini et al. 2015, 2017a, b), as a part of the revisionary study on the genus *Argyreia* in India. In this paper, names of two more species are lectotypified in accordance with Article 9.3 of the the International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code. Turland et al. 2018). Details of the types for each taxon, distribution (worldwide and in India) and other required notes wherever necessary are provided.

1. *Argyreia fulgens* Choisy in Mém. Soc. Phys. Genève **6**: 415 (1834) & Convolv. Orient.: 33 (1834).

Lectotype (designated here): INDIA. Without locality, 3 September 18 [14?], Wallich Numer. List No. 1394. 1 (K001113007, specimen on the left, Fig. 1; K000830711–isolectotype).

Distribution: Endemic to India (Kerala and

Tamil Nadu).

In the protologue, Choisy (1834) states that the type is based on Wallich's Numerical List (also known as Wallich's Catalogue) entry 1394 that belonged to Heyne's herbarium and Madras Herbarium. There is one herbarium sheet at K-W which has two specimens with original labels. The specimen on the left side corresponding to Wallich 1394.1 (K001113007), collected on 3 September 18[14?], has been chosen as the lectotype for the name as it is well-preserved with flowers and matches well with the description given in the protologue. The one on the right side corresponds to Wallich 1394B (K001113008) was collected on October 1814. Apart, there are three specimens one each at K (K000830711), CAL (Acc. No. 310680) and G-DC (G00135017) with the same Wallich's Numerical List entry 1394. Among these, the specimen with barcode K000830711 is considered as isolectotype.

2. *Argyreia lanceolata* Choisy in Mém. Soc. Phys. Genève **6**: 421 (1834) & Convolv. Orient.: 39 (1834).

Lectotype (designated here): MYANMAR. Tavoy, 2 Oct. 1827, W.Gomez 349, Wallich Numer. List No. 1395.2 (K-W, K001113009, Fig. 2; G-DC, G00134913–isolectotype).

Distribution: India (Andaman and Nicobar

as the lectotype while the duplicate specimen at G-DC, is considered as isolectotype.

The authors are grateful to the Director, Botanical Survey of India and Head of Office, Central National Herbarium, Howrah, for facilities and encouragement.

References

- Choisy J. 1834. *Convolvulaceae* Orientale. Mém. Soc. Phys. Genève **6**(2): 383–502.
- Shalini S., Lakshminarasimhan P. and Arisdason W. 2015. Lectotypification of *Argyreia bella* (*Convolvulaceae*). *Telopea* **18**: 391–393.
- Shalini S., Lakshminarasimhan P. and Arisdason W. 2017a. Lectotypification of four names in *Argyreia* (*Convolvulaceae*). *Telopea* **20**: 171–177.
- Shalini S., Lakshminarasimhan P. and Maity D. 2017b. A new combination and lectotypification in *Argyreia* (*Convolvulaceae*). *Nelumbo* **59**(2): 145–148.
- Turland N.J., Wiersema J.H., Barrie F.R., Greuter W., Hawksworth D.L., Herendeen P.S., Knapp S., Kusber W.H., Li D.-Z., Marhold K., May T.W., McNeill J., Monro A.M., Prado J., Price M.J. and Smith G.F. 2018. International Code of Nomenclature for algae, fungi and plants (Shenzhen Code). *Regnum Vegetabile* 159. Koeltz Scientific Books, Koenigstein. <https://doi.org/10.12705/Code.2018>.
- Wallich N. 1824. A Numerical List of Dried Specimens of Plants in the East India Company's Museum: Collected under the Superintendence of Dr. Wallich of the Company's Botanic Garden at Calcutta. London.

S.Shalini^a, P.Lakshminarasimhan^b, W.Arisdason^c, D.Maity^d: ヒルガオ科 *Argyreia* 2種のレクトタイプ指定
インド産の *Argyreia* Lour.(ヒルガオ科ギンヨウアサガオ属) の2種 *A. fulgens* Choisy (Fig. 1) および *A. lanceolata* Choisy (Fig. 2) のレクトタイプを指定した。
(^aインド・Botanical Survey of India,

Central National Herbarium,
^bインド・Uttam Townscapes, Maharashtra,
^cインド・Botanical Survey of India, Southern Regional Centre,
^dインド・University of Calcutta, Department of Botany)