

○*Nardia* 属の分類システム (井上 浩) Hiroshi INOUE: Nomenclatural changes of subgeneric and sectional ranks of the genus *Nardia* Gray

植研 46 (1): 1-7 (1971) に私はシッキムで採集された *Nardia flagelliformis* を新種として記載すると共に, *Nardia* 属の新らしい分類システムを提唱した。このシステムを提唱するにあたって, この属の基準種を *Nardia scalaris* (Schrad.) Gray と考える Evans や Schuster の考えを採って, subgenus ならびに section の命名をおこなった。然し, *Nardia* 属の基準種としては *N. compressa* (Hook.) Gray を用いる方が正しいことが判明したので, 以下の通り訂正する。

Genus *Nardia* Gray, Nat. Arr. Brit. Pl. 1: 694 (1821). Type species: *Nardia compressa* (Hook.) Gray.

Subg. *Nardia* Syn. *Nardia* sect. *Compressae* Schust. (1969) = *Nardia* sect. *Mesophylla* (Dum.) Carr. (1874) = *Nardia* subg. *Mesophylla* (Dum.) Lindb., Meddel. Soc. Fauna Fl. Fenn. 13: 238 (1886); Inoue, Journ. Jap. Bot. 46: 6 (1971).

Subg. *Geoscypharia* Trev., Mem. Real. Institut. Lombard. 13: 401 (1877). Type species: *Nardia geoscyphos* (DeNot.) Lindb. Syn. *Nardia* subg. *Gongylothalamus* Schiffn. (1898).

The present subgenus includes the following four sections.

Sect. *Geoscypharia* This section includes the species listed under subg. *Nardia* sect. *Nardia* by Inoue (Journ. Jap. Bot. 41: 6. 1971).

Sect. *Subclavatae* (Inoue) Inoue, comb. nov. = *Nardia* Subgen. *Nardia* Sect. *Subclavatae* Inoue, Journ. Jap. Bot. 46: 6 (1971). Type species: *Nardia subclavata* (Steph.) Amak.

Sect. *Breidlerion* Grolle, Bot. Mag. Tokyo 77: 299 (1964). Type species: *Nardia breidlerii* (Limpr.) Lindb.

Sect. *Flagelliformes* (Inoue) Inoue, comb. nov. = *Nardia* subg. *Nardia* sect. *Flagelliformes* Inoue, Journ. Jap. Bot. 46: 6 (1971). Type species: *Nardia flagelliformis* Inoue.

The subgeneric and sectional nomenclatures were changed from my previous proposal (Inoue, in Journ. Jap. Bot. 46: 1-7. 1971). Although Schuster (The Hepaticae and Anthocerotae of N. America, vol. 2, 1969) accepted *Nardia scalaris* (Schrad.) Gray as the type species of the genus, *N. compressa* (Hook.) Gray should be the type (*conserved*!).

(国立科学博物館)